

# A stability version for a theorem of Erdős on nonhamiltonian graphs

Zoltán Füredi\*

Alexandr Kostochka†

Ruth Luo‡

August 23, 2016

## Abstract

Let  $n, d$  be integers with  $1 \leq d \leq \lfloor \frac{n-1}{2} \rfloor$ , and set  $h(n, d) := \binom{n-d}{2} + d^2$  and  $e(n, d) := \max\{h(n, d), h(n, \lfloor \frac{n-1}{2} \rfloor)\}$ . Because  $h(n, d)$  is quadratic in  $d$ , there exists a  $d_0(n) = (n/6) + O(1)$  such that

$$e(n, 1) > e(n, 2) > \cdots > e(n, d_0) = e(n, d_0 + 1) = \cdots = e(n, \lfloor \frac{n-1}{2} \rfloor).$$

A theorem by Erdős states that for  $d \leq \lfloor \frac{n-1}{2} \rfloor$ , any  $n$ -vertex nonhamiltonian graph  $G$  with minimum degree  $\delta(G) \geq d$  has at most  $e(n, d)$  edges, and for  $d \geq d_0(n)$  the unique sharpness example is simply the graph  $K_n - E(K_{\lceil (n+1)/2 \rceil})$ . Erdős also presented a sharpness example  $H_{n,d}$  for each  $1 \leq d \leq d_0(n)$ .

We show that if  $d < d_0(n)$  and a 2-connected, nonhamiltonian  $n$ -vertex graph  $G$  with  $\delta(G) \geq d$  has more than  $e(n, d+1)$  edges, then  $G$  is a subgraph of  $H_{n,d}$ . Note that  $e(n, d) - e(n, d+1) = n - 3d - 2 \geq n/2$  whenever  $d < d_0(n) - 1$ .

**Mathematics Subject Classification:** 05C35, 05C38.

**Keywords:** Turán problem, hamiltonian cycles, extremal graph theory.

Dedicated to the memory of Professor H. Sachs.

## 1 Introduction

We use standard notation. In particular,  $V(G)$  denotes the vertex set of a graph  $G$ ,  $E(G)$  denotes the edge set of  $G$ , and  $e(G) = |E(G)|$ . Also, if  $v \in V(G)$ , then  $N(v)$  denotes the neighborhood of  $v$  and  $d(v) = |N(v)|$ . Ore [4] proved the following Turán-type result:

**Theorem 1** (Ore [4]). *If  $G$  is a nonhamiltonian graph on  $n$  vertices, then  $e(G) \leq \binom{n-1}{2} + 1$ .*

This bound is achieved only for the  $n$ -vertex graph obtained from the complete graph  $K_{n-1}$  by adding a vertex of degree 1. Erdős [2] refined the bound in terms of the minimum degree of the graph:

---

\*Alfréd Rényi Institute of Mathematics, Hungary E-mail: [furedi.zoltan@renyi.mta.hu](mailto:furedi.zoltan@renyi.mta.hu). Research supported in part by the Hungarian National Science Foundation OTKA 104343, by the Simons Foundation Collaboration Grant 317487, and by the European Research Council Advanced Investigators Grant 267195.

†University of Illinois at Urbana-Champaign, Urbana, IL 61801 and Sobolev Institute of Mathematics, Novosibirsk 630090, Russia. E-mail: [kostochk@math.uiuc.edu](mailto:kostochk@math.uiuc.edu). Research of this author is supported in part by NSF grants DMS-1266016 and DMS-1600592 and grants 15-01-05867 and 16-01-00499 of the Russian Foundation for Basic Research.

‡University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA. E-mail: [ruthluo2@illinois.edu](mailto:ruthluo2@illinois.edu).

**Theorem 2** (Erdős [2]). *Let  $n, d$  be integers with  $1 \leq d \leq \lfloor \frac{n-1}{2} \rfloor$ , and set  $h(n, d) := \binom{n-d}{2} + d^2$ . If  $G$  is a nonhamiltonian graph on  $n$  vertices with minimum degree  $\delta(G) \geq d$ , then*

$$e(G) \leq \max \left\{ h(n, d), h(n, \left\lfloor \frac{n-1}{2} \right\rfloor) \right\} =: e(n, d).$$

*This bound is sharp for all  $1 \leq d \leq \lfloor \frac{n-1}{2} \rfloor$ .*

To show the sharpness of the bound, for  $n, d \in \mathbb{N}$  with  $d \leq \lfloor \frac{n-1}{2} \rfloor$ , consider the graph  $H_{n,d}$  obtained from a copy of  $K_{n-d}$ , say with vertex set  $A$ , by adding  $d$  vertices of degree  $d$  each of which is adjacent to the same  $d$  vertices in  $A$ . An example of  $H_{11,3}$  is below.




Figure 1:  $H_{11,3}$

By construction,  $H_{n,d}$  has minimum degree  $d$ , is nonhamiltonian, and  $e(H_{n,d}) = \binom{n-d}{2} + d^2 = h(n, d)$ . Elementary calculation shows that  $h(n, d) > h(n, \lfloor \frac{n-1}{2} \rfloor)$  in the range  $1 \leq d \leq \lfloor \frac{n-1}{2} \rfloor$  if and only if  $d < (n+1)/6$  and  $n$  is odd or  $d < (n+4)/6$  and  $n$  is even. Hence there exists a  $d_0 := d_0(n)$  such that

$$e(n, 1) > e(n, 2) > \cdots > e(n, d_0) = e(n, d_0 + 1) = \cdots = e(n, \left\lfloor \frac{n-1}{2} \right\rfloor),$$

where  $d_0(n) := \lceil \frac{n+1}{6} \rceil$  if  $n$  is odd, and  $d_0(n) := \lceil \frac{n+4}{6} \rceil$  if  $n$  is even. Let  $H'_{n,d}$  denote the graph that is an edge-disjoint union of two complete graphs  $K_{n-d}$  and  $K_{d+1}$  sharing one vertex.

The result of this note is the following refinement of Theorem 2.

**Theorem 3.** *Let  $n \geq 3$  and  $d \leq \lfloor \frac{n-1}{2} \rfloor$ . Suppose that  $G$  is an  $n$ -vertex nonhamiltonian graph with minimum degree  $\delta(G) \geq d$  such that*

$$e(G) > e(n, d+1) = \max \left\{ h(n, d+1), h(n, \left\lfloor \frac{n-1}{2} \right\rfloor) \right\}. \quad (1)$$

*(So we have  $d < d_0(n)$ .) Then  $G$  is a subgraph of either  $H_{n,d}$  or  $H'_{n,d}$ .*

This is a stability result in the sense that for  $d < n/6$ , each 2-connected, nonhamiltonian  $n$ -vertex graph with minimum degree at least  $d$  and “close” to  $h(n, d)$  edges is a subgraph of the extremal graph  $H_{n,d}$ . Note that  $h(n, d) - h(n, d+1) = n - 3d - 2$  is at least  $n/2$  for  $d < d_0 - 1$ . Note also that  $e(H'_{n,d}) > e(n, d+1)$  only when  $d = O(\sqrt{n})$ .

We will use the following well-known theorems of Pósa.

**Theorem 4** (Pósa [5]). *Let  $n \geq 3$ . If  $G$  is a nonhamiltonian  $n$ -vertex graph, then there exists  $1 \leq k \leq \lfloor \frac{n-1}{2} \rfloor$  such that  $G$  has a set of  $k$  vertices with degree at most  $k$ .*

**Theorem 5** (Pósa [6]). *Let  $n \geq 3$ ,  $1 \leq \ell < n$  and let  $G$  be an  $n$ -vertex graph such that  $d(u) + d(v) \geq n + \ell$  for every non-edge  $uv$  in  $G$ . Then for every linear forest  $F$  with  $\ell$  edges contained in  $G$ , the graph  $G$  has a hamiltonian cycle containing all edges of  $F$ .*

## 2 Proof of Theorem 3

Call a graph  $G$  *saturated* if  $G$  is nonhamiltonian but for each  $uv \notin E(G)$ ,  $G + uv$  has a hamiltonian cycle. Ore's proof [4] of Dirac's Theorem [1] yields that

$$\text{for every } n\text{-vertex saturated graph } G \text{ and for each } uv \notin E(G), d(u) + d(v) \leq n - 1. \quad (2)$$

First we show two facts on saturated graphs with many edges.

**Lemma 6.** *Let  $G$  be a saturated  $n$ -vertex graph with  $e(G) > h(n, \lfloor \frac{n-1}{2} \rfloor)$ . Then for some  $1 \leq k \leq \lfloor \frac{n-1}{2} \rfloor$ ,  $V(G)$  contains a subset  $D$  of  $k$  vertices of degree at most  $k$  such that  $G - D$  is a complete graph.*

*Proof.* Since  $G$  is nonhamiltonian, by Theorem 4, there exists some  $1 \leq k \leq \lfloor \frac{n-1}{2} \rfloor$  such that  $G$  has  $k$  vertices with degree at most  $k$ . Pick the maximum such  $k$ , and let  $D$  be the set of the vertices with degree at most  $k$ . Since  $e(G) > h(n, \lfloor \frac{n-1}{2} \rfloor)$ ,  $k < \lfloor \frac{n-1}{2} \rfloor$ . So, by the maximality of  $k$ ,  $|D| = k$ .

Suppose there exist  $x, y \in V(G) - D$  such that  $xy \notin E(G)$ . Among all such pairs, choose  $x$  and  $y$  with the maximum  $d(x)$ . Since  $y \notin D$ ,  $d(y) > k$ . Let  $D' := V(G) - N(x) - \{x\}$  and  $k' := |D'| = n - 1 - d(x)$ . By (2),

$$d(z) \leq n - 1 - d(x) = k' \text{ for all } z \in D'. \quad (3)$$

So  $D'$  is a set of  $k'$  vertices of degree at most  $k'$ . Since  $y \in D'$ ,  $k' \geq d(y) > k$ . Thus by the maximality of  $k$ , we get  $k' = n - 1 - d(x) > \lfloor \frac{n-1}{2} \rfloor$ . Equivalently,  $d(x) < \lceil \frac{n-1}{2} \rceil$ . For all  $z \in D' + \{x\}$ , either  $z \in D$  where  $d(z) \leq k \leq \lfloor \frac{n-1}{2} \rfloor$ , or  $z \in V(G) - D$ , and so  $d(z) \leq d(x) \leq \lfloor \frac{n-1}{2} \rfloor$ . It follows that  $e(G) \leq h(n, \lfloor \frac{n-1}{2} \rfloor)$ , a contradiction.  $\square$

**Lemma 7.** *Under the conditions of Lemma 6, if  $k = \delta(G)$ , then  $G = H_{n, \delta(G)}$  or  $G = H'_{n, \delta(G)}$ .*

*Proof.* Set  $d := \delta(G)$ , and let  $D$  be a set of  $d$  vertices with degree at most  $d$ . Let  $u \in D$ . Since  $\delta(G) \geq |D| = d$ ,  $u$  has a neighbor  $w \in V(G) - D$ . Consider any  $v \in D - \{u\}$ . By Lemma 6,  $w$  is adjacent to all of  $V(G) - D - \{w\}$ . It also is adjacent to  $u$ , therefore its degree is at least  $n - d$ . We obtain

$$d(w) + d(v) \geq (n - d) + d = n.$$

Then by (2),  $w$  is adjacent to  $v$ , and hence  $w$  is adjacent to all vertices of  $D$ .

Let  $W$  be the set of vertices in  $V(G) - D$  having a neighbor in  $D$ . We have obtained that  $W \neq \emptyset$  and

$$N(u) \cap (V(G) - D) = W \text{ for all } u \in D. \quad (4)$$

Let  $G' = G[D \cup W]$ . If  $|W| = 1$ , then  $G = H'_{n,d}$ . If  $|V(G')| = 2d$ , then by (4), each vertex  $u \in D$  has the same  $d$  neighbors in  $V(G) - D$ . Because  $d(u) = d$ ,  $D$  is an independent set. Thus  $G = H_{n,d}$ . Otherwise,  $d + 2 \leq |V(G')| \leq 2d - 1$ ,  $|D| \geq 2$ .

Fix a pair of vertices  $w_1, w_2 \in W$ . For any  $x, y \in V(G')$ ,

$$d(x) + d(y) \geq d + d \geq |V(G')| + 1.$$

Therefore by Theorem 5,  $G'$  has a hamiltonian cycle  $C$  that uses the edge  $w_1w_2$ . Since  $G'' := G - (V(G') - \{w_1, w_2\})$  is a complete graph, it contains a hamiltonian  $w_1, w_2$ -path  $P$ . Then  $P \cup (C - w_1w_2)$  is a hamiltonian cycle of  $G$ , a contradiction.  $\square$

*Proof of Theorem 3.* Suppose that an  $n$ -vertex, nonhamiltonian graph  $G$  satisfies the constraints of Theorem 3 for some  $1 \leq d \leq \lfloor \frac{n-1}{2} \rfloor$ . We may assume  $G$  is saturated, since if a graph containing  $G$  is a subgraph of  $H_{n,d}$  or  $H'_{n,d}$ , then  $G$  is as well.

By Lemma 6,  $G$  has a set  $D$  of  $k \leq \lfloor \frac{n-1}{2} \rfloor$  vertices with degree at most  $k$  such that  $G - D$  is a complete graph. Therefore  $e(G) \leq \binom{n-k}{2} + k^2 = h(n, k)$ . If  $k > d$ , then  $e(G) \leq \max\{h(n, d + 1), h(n, \lfloor \frac{n-1}{2} \rfloor)\} = e(n, d + 1)$ , a contradiction. Thus  $k \leq d$ . Furthermore,  $k \geq \delta(G) \geq d$ , and hence  $k = d$ . Also, since  $e(G) > h(n, \lfloor \frac{n-1}{2} \rfloor)$ , we have  $d + 1 \leq d_0(n) \leq (n + 8)/6$ . Applying Lemma 7 completes the proof.  $\square$

**Acknowledgment.** We thank both referees for their helpful comments.

**Acknowledgment added on April 5, 2017.** We have learned that Theorem 3 has already been proved by Li and Ning as a lemma in [3] with a somewhat different proof.

## References

- [1] G. Dirac, Some theorems on abstract graphs, Proc. London Math. Soc. (3) **2** (1952), 69–81.
- [2] P. Erdős, Remarks on a paper of Pósa, Magyar Tud. Akad. Mat. Kutató Int. Közl. **7** (1962), 227–229.
- [3] B. Li and B. Ning, Spectral analogues of Erdős’ and Moon-Moser’s theorems on Hamilton cycles, Linear Multilinear Algebra, **64** (2016), no.11, 1152–1169.
- [4] O. Ore, Arc coverings of graphs, Acta Math. Acad. Sci. Hung. **10** (1959), 337–356.
- [5] L. Pósa, A theorem concerning Hamilton lines, Magyar Tud. Akad. Mat. Kutató Int. Közl. **7** (1962), 225–226.
- [6] L. Pósa, On the circuits of finite graphs, Magyar. Tud. Akad. Mat. Kutató Int. Közl. **8** (1963/1964), 355–361.